

HYGRO BOOST-FLO

MICRO-NUTRIENT MIXTURE / MIKRO-ELEMENT MENGSEL

Nitrogen / Stikstof (N)	198 g/L	152 g/kg
Manganese / Mangaan (Mn)	20.7 g/L	15.9 g/kg
Zinc / Sink (Zn)	50.0 g/L	38.5 g/kg
Copper / Koper (Cu)	32.7 g/L	25.2 g/kg
Boron / Boor (B)	8.7 g/L	6.7 g/kg

SG: 1.3 @ 20°C

FERTILIZER GRP 1 MISSTOF

Reg. No. / Nr. K6252

Act No. / Wet Nr. 36 of / van 1947

DIRECTIONS FOR USE

Use as directed in the following table.

Always use in combination with Nu-Film P®* or Nu-Film 17®*.

Shake well before use.

*Nu-Film P® L2980 Act No. 36 of 1947

*Nu-Film 17® L2981 Act No. 36 of 1947

PRECAUTIONS

Store in a well-sealed container away from direct sunlight.

Keep away from children, uninformed persons, animals and feed.

Wash hands after use.

APPLICATION QUANTITIES

CROPS	RATE	RECOMMENDATIONS
Young non-bearing citrus trees	2-3 L/ha	Apply at 4-6 week intervals during the months of August to February. Apply as a full cover spray.
Fruit trees	2-3 L/ha	Apply at 6 week intervals during periods of active vegetative growth. Especially suited for application on young trees. Apply as a full cover spray.
Ornamental trees	2-3 L/ha	Apply at regular intervals during vegetative growth or as a corrective spray. It is also recommended as a post-transplant spray to initiate growth. Apply as a full cover spray.
Woody shrubs	2-3 L/ha	Apply at regular intervals during vegetative growth or as a corrective spray. It is also recommended as a post-transplant spray to initiate growth. Apply as a full cover spray.
Field crops i.e. wheat, beans, maize etc.	2-3 L/ha	Apply as a full cover spray during period of active growth.
Vegetable crops	2-3 L/ha	Apply as a full cover spray during period of active growth.

GEBRUIKSAANWYSINGS

Gebruik soos aangedui in bygaande tabel.

Gebruik altyd saam met Nu-Film P®* of Nu-Film 17®*.

Skud goed voor gebruik.

*Nu-Film P® L2980 Wet Nr. 36 van 1947

*Nu-Film 17® L2981 Wet Nr. 36 van 1947

VOORSORGMAATREËLS

Bêre in 'n verseëelde houer weg van direkte sonlig.

Hou weg van kinders, oningeligte persone, diere en voer.

Was hande na gebruik.

TOEDIENINGSHOEVEELHEDE

GEWAS	DOSIS	AANBEVELINGS
Jong nie-draende sitrus bome	2-3 L/ha	Dien toe met 4-6 weke intervalle tussen Augustus en Februarie. Dien toe as 'n algehele bedekkingspuit.
Vrugte bome	2-3 L/ha	Dien toe met 6 weke intervalle tydens periodes van aktiewe vegetatiewe groei. Veral geskik vir toediening op jong bome. Dien toe as 'n algehele bedekkingspuit.
Ornamentele bome	2-3 L/ha	Dien toe met gereelde intervalle tydens periodes van aktiewe vegetatiewe groei of as 'n korrektiewe spuit. Word ook aanbeveel as 'n na-uitplant toediening om aktiewe groei te bevorder. Dien toe as 'n algehele bedekkingspuit.
Houtagtige struik	2-3 L/ha	Dien toe met gereelde intervalle tydens periodes van aktiewe vegetatiewe groei of as 'n korrektiewe spuit. Word ook aanbeveel as 'n na-uitplant toediening om aktiewe groei te bevorder. Dien toe as 'n algehele bedekkingspuit.
Akkerbou gewasse bv. koring, bone, mielies ens.	2-3 L/ha	Dien toe as 'n algehele bedekkingspuit tydens periodes van aktiewe groei.
Groente gewasse	2-3 L/ha	Dien toe as 'n algehele bedekkingspuit tydens periodes van aktiewe groei.

Manufactured, Registered & Distributed by / Vervaardig, Gereistreer & Versprei deur:



PROPERTIES (PTY) LTD

Reg. No. 1984/00638/07

1 Gerard Braak St, Pyramid

PO Box 17220, Pretoria North, 0116

Tel (012) 545 8000 • Fax (012) 545 8088