

HYGROPLEX PLUS

Reg. No. **B 5594** Act No. 36 of 1947

Fertilizer Group 2

Water soluble EDTA chelated micro nutrient mixture

Iron (Fe)*	84 g/kg
Manganese (Mn)*	20 g/kg
Zinc (Zn)*	10 g/kg
Copper (Cu)*	1.5 g/kg
Boron (B)	25 g/kg
Molybdenum (Mo)	3 g/kg

*EDTA Chelated

DIRECTION FOR USE: Use only as directed

HYGROPLEX PLUS is recommended for use during plant growth periods that require relatively high micro-nutrient levels and for the correction of trace element deficiencies on crops by means of soil application and / or foliar application.

PRECAUTIONS

- Store in a well-sealed container away from light.
- Keep away from children, uninformed persons, animals and feedstuff.
- Wash hands after use.

APPLICATION RATES

Crops	Rate	Recommendations
Pot plants and shrubs (Hydroponic)	10-20 g /1000 ℓ	Apply continuously in the feeding system.
Greenhouse cucumbers and tomatoes (Hydroponics)	20 g /1000 ℓ	Apply in the feeding solution 2-3 times per growing season. The final solution will contain the following nutrient levels in mg/ ℓ (ppm): Fe –1.68 Mn – 0.4 Zn–0.2 Cu –0.03 Mo – 0.06
Seedlings	10-20 g /1000 ℓ	Apply 2-3 times in the feeding solution from emergence to transplant,
Agricultural crops and vegetable crops	1-3 kg / ha	Foliar application: Apply 0.5 -0.15 kg/ha or 1-2 g/ℓ as early in the season as possible, using a fine spray nozzle. Do not exceed spray concentrations of 0.2%. Where severe deficiencies occur, repeat application 2-3 times at 7-10 day intervals. Soil Application: Dissolve powder in a convenient volume of water e.g. (10 g/ℓ). Apply as a coarse spray over the root zone. Alternatively apply as powder to the root zone but incorporate immediately into top soil by irrigation.

HYGROPLEX PLUS

Reg. Nr. **B 5594** Wet Nr. 36 van 1947

Groep 2 Misstof

'n water oplosbare EDTA gecheleerde mikro-element mengsel

Yster (Fe)*	84 g/kg
Mangaan (Mn)*	20 g/kg
Sink (Zn)*	10 g/kg
Koper (Cu)*	1.5 g/kg
Boor (B)*	25 g/kg
Molibdeen (Mo)	3 g/kg

* EDTA gecheleerde

GEBRUIKSAANWYSINGS: Gebruik slegs soos aangedui

HYGROPLEX PLUS word aanbeveel vir gebruik gedurende plantgroeitydperke wat relatief hoë mikrovoedingstowwe benodig en vir die regstelling van tekorte aanmikro-elemente in gewasse deur middel van grondtoediening en / of blaartoediening.

VOORSOMAGMAATREËLS

- Bêre in 'n goed verseëlde houer weg van lig.
- Hou weg van kinders, oningeligte persone, diere en voere.
- Was hande ná gebruik.

TOEDIENINGS HOEEVEELHEDE

Gewasse	Dosis	Aanbevelings
Potplante en struik (Hydroponies)	10-20 g /1000 ℓ	Toegedien voortdurend in die voedingsstelsel toegepas.
Komkommers en tamaties in kweekhuise (Hydroponies)	20 g /1000 ℓ	Toegedien in die voedingsoplossing 2-3 keer per groeiseisoen. Die finale oplossing sal die volgende voedingsvlakke in mg / ℓ (ppm) bevat: Fe –1.68 Mn – 0.4 Zn–0.2 Cu –0.03 Mo – 0.06
Saailinge	10-20 g /1000 ℓ	Toegedien in die voedingsoplossing 2-3 keer per groei seisoen,
Landbougewasse en groente	1-3 kg / ha	Blaarvoeding: Spuut 0.5 - 1.5 kg/ha of 1-2 g/ℓ so vroeg moontlik in die seisoen, met 'n fyn spuit nevel. Moenie die spuitkonsentrasie van 0.2 % oorskry nie. Waarernstige tekorte voorkom, herhaal 2-3 keer met 7-10 dae tussenposes. Grondtoediening: Los poeier op in gerieflike hoeveelheid water bv. (10 g/ℓ). Dien toe oor die wortel gebied deur gebruik te maak van 'n growwe sproei. Alternatiewelik dien as poeier direk op grond toe, maar inkorporeer onmiddellik in die boonste grondlae d.m.v besproeiing.

Manufactured, Registered and Distributed by / Vervaardig, Geregistreer en Versprei deur:



PROPERTIES (PTY) LTD. • Reg. No. 1984/00638/07

1 Gerard Braak Street, Pyramid, 0120

P.O Box 17220, Pretoria North, 0116

Tel: (012) 545 8000 • Fax: (012) 545 8088